

# Prophylaxis against infective endocarditis: antimicrobial prophylaxis against infective endocarditis in adults and children undergoing interventional procedures

Clinical guideline

Published: 17 March 2008

[nice.org.uk/guidance/cg64](https://www.nice.org.uk/guidance/cg64)

## Contents

Recommendations .....	3
Adults and children with structural cardiac defects at risk of developing infective endocarditis.....	3
Patient advice .....	3
Prophylaxis against infective endocarditis .....	4
Infection .....	4
Context.....	6
Recommendations for research .....	7
1 National register of infective endocarditis .....	7
2 Cardiac conditions and infective endocarditis .....	7
3 Interventional procedures and infective endocarditis.....	7
4 Antibiotic prophylaxis against infective endocarditis .....	8
Update information .....	9

## Recommendations

People have the right to be involved in discussions and make informed decisions about their care, as described in [your care](#).

[Using NICE guidelines to make decisions](#) explains how we use words to show the strength of our recommendations, and has information about safeguarding, consent and prescribing medicines (including 'off-label' use).

### *Adults and children with structural cardiac defects at risk of developing infective endocarditis*

1.1.1 Healthcare professionals should regard people with the following cardiac conditions as being at risk of developing infective endocarditis:

- acquired valvular heart disease with stenosis or regurgitation
- hypertrophic cardiomyopathy
- previous infective endocarditis
- structural congenital heart disease, including surgically corrected or palliated structural conditions, but excluding isolated atrial septal defect, fully repaired ventricular septal defect or fully repaired patent ductus arteriosus, and closure devices that are judged to be endothelialised
- valve replacement. [2015]

### *Patient advice*

1.1.2 Healthcare professionals should offer people at risk of infective endocarditis clear and consistent information about prevention, including:

- the benefits and risks of antibiotic prophylaxis, and an explanation of why antibiotic prophylaxis is no longer routinely recommended
- the importance of maintaining good oral health
- symptoms that may indicate infective endocarditis and when to seek expert advice

- the risks of undergoing invasive procedures, including non-medical procedures such as body piercing or tattooing. [2015]

## *Prophylaxis against infective endocarditis*

1.1.3 Antibiotic prophylaxis against infective endocarditis is not recommended:

- for people undergoing dental procedures
- for people undergoing non-dental procedures at the following sites<sup>[1]</sup>:
  - upper and lower gastrointestinal tract
  - genitourinary tract; this includes urological, gynaecological and obstetric procedures, and childbirth
  - upper and lower respiratory tract; this includes ear, nose and throat procedures and bronchoscopy. [2015]

1.1.4 Chlorhexidine mouthwash should not be offered as prophylaxis against infective endocarditis to people at risk of infective endocarditis undergoing dental procedures. [2015]

## *Infection*

1.1.5 Any episodes of infection in people at risk of infective endocarditis should be investigated and treated promptly to reduce the risk of endocarditis developing. [2015]

1.1.6 If a person at risk of infective endocarditis is receiving antimicrobial therapy because they are undergoing a gastrointestinal or genitourinary procedure at a site where there is a suspected infection, the person should receive an antibiotic that covers organisms that cause infective endocarditis. [2015]

You can also see this guideline in the NICE pathway on [prophylaxis against infective endocarditis](#).

To find out what NICE has said on topics related to this guideline, see our web page on [cardiovascular conditions: general and other](#).

<sup>[1]</sup>The evidence reviews for this guideline covered only procedures at the sites listed in this recommendation. Procedures at other sites are outside the scope of the guideline (see the [scope](#) for details).

## Context

Infective endocarditis is a rare condition with significant morbidity and mortality. It may arise after bacteraemia in a person with a predisposing cardiac lesion.

In the past, people at risk of infective endocarditis were given antibiotic prophylaxis before dental and certain non-dental interventional procedures. However, the British Society for Antimicrobial Chemotherapy [Guidelines for the prevention of endocarditis](#) (2006) and the American Heart Association guideline [Prevention of infective endocarditis](#) (2007) recommended changes in clinical practice that aimed to limit the use of antibiotic prophylaxis in people having these procedures.

In 2008 NICE published a guideline to give clear, evidence-based guidance on best clinical practice for prophylaxis against infective endocarditis in people having dental and certain non-dental interventional procedures. The guideline recommended that people at risk of infective endocarditis having interventional procedures (dental procedures, upper and lower respiratory tract procedures, upper and lower gastrointestinal tract procedures, genitourinary tract procedures) should not be offered antibiotic prophylaxis against infective endocarditis. It also recommended which patients with pre-existing cardiac lesions should be regarded as at risk. These recommendations marked a further change from accepted practice.

The incidence of infective endocarditis has been shown to be increasing over time. The reasons for this are not well understood. However, in 2014 the Lancet published a paper that reported an increase in cases of infective endocarditis from 2000 to 2013, which showed a higher rate of increase following the publication of the NICE guideline ([Incidence of infective endocarditis in England, 2000–13](#) Dayer MJ et al). In light of this paper, NICE felt it was important to assess any new evidence published since the 2008 NICE guideline. This evidence has been taken into account in the 2015 update of this guideline.

## Recommendations for research

In 2008 the Guideline Committee made 3 recommendations for research.

As part of the 2015 update, the Standing Committee made an additional research recommendation on antibiotic prophylaxis against infective endocarditis. This can be found in the [addendum](#).

### *1 National register of infective endocarditis*

Infective endocarditis is a rare condition. The development of a national register of infective endocarditis in the UK to support research is recommended.

#### **Why this is important**

Such research would be facilitated by the availability of a national register that could offer data into the 'case' arm of proposed case-control studies and should be an anonymised database that would not require patient consent and hence more straightforward case ascertainment. Although it is a rare condition, it is likely that across the country there are enough patients to generate evidence from well conducted national studies.

### *2 Cardiac conditions and infective endocarditis*

What is the risk of developing infective endocarditis in people with acquired valvular disease and structural congenital heart disease?

#### **Why this is important**

Such research should use a population-based cohort study design to allow direct comparison between groups and allow estimation of both relative and absolute risk.

### *3 Interventional procedures and infective endocarditis*

What is the frequency and level of bacteraemia caused by non-oral daily activities (for example, urination or defaecation)?

#### **Why this is important**

Such research should quantitatively determine the frequency and level of bacteraemia.

## ***4 Antibiotic prophylaxis against infective endocarditis***

Does antibiotic prophylaxis in those at risk of developing infective endocarditis reduce the incidence of infective endocarditis when given before a defined interventional procedure?

### **Why this is important**

There is limited evidence about the effectiveness of antibiotic prophylaxis in reducing the incidence of infective endocarditis in people at risk of developing infective endocarditis. The current evidence includes very limited data from observational studies with inconclusive findings. The study should be a randomised controlled trial with long-term follow-up comparing antibiotic prophylaxis with no antibiotic prophylaxis in adults and children with underlying structural cardiac defects undergoing interventional procedures. Outcomes should include the incidence infective endocarditis in those receiving prophylaxis compared to those not, and the incidence of adverse effects including anaphylaxis.



## Update information

**September 2015:** This guideline is an update of NICE guideline CG64 (published March 2008). No new recommendations have been added. For recommendations marked as [2015] the evidence has been reviewed but no change has been made to the recommended action.

ISBN: 978-1-4731-1434-0

## Accreditation

